



Estimating Wheat Grain Yield Potential

Jeff Edwards
Small Grains Extension Specialist

Wheat grain yield is the product of heads per square foot, seeds per head and seed weight. Using estimates of these parameters, farmers can derive an estimated grain yield potential. These estimates are generally more accurate as the season progresses. An estimate of the number of heads per square foot in the late spring, for example, will be much more accurate than one made in the fall. It is also important to note that any estimate is an *estimate*, and nothing more, until the crop is harvested.

Heads per Square Foot


Heads per square foot is dictated by seeding rate, tillering and tiller survival. Most wheat seed in Oklahoma has between 13,000 and 16,000 seeds per pound. At a 60 lbs/A seeding rate, this will deliver 18 seeds to 22 seeds per square foot. Wheat needs 60 heads to 70 heads per square foot at harvest for full yield potential. October-sown wheat will generally produce 2 to 3 good tillers per plant and will reach the 60 heads per square foot target when sown at 60 lbs/A. November-sown wheat may have only 1 to 2 good tillers per plant and will, therefore, fall short of the 60 heads per square foot target when sown at 60 lbs/A.

Potential heads per square foot can be estimated in field by counting tillers in one linear foot of row and using Equation 1 to calculate the number of tillers per square foot. These calculations have been performed for various row spacings and tiller counts in Table 1. It is important to count only tillers with at least three unfolded leaves. It is also important to keep in mind that these are tillers that *can* produce grain heads; environmental conditions could still cause the plant to abort some of these tillers. The closer to harvest that these counts are made, the more accurate the estimate will be.

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: <http://osufacts.okstate.edu>

Table 1. Tillers per square foot based on tiller counts per linear foot of row at various row spacings.

Tillers per foot of row	Row Spacing in Inches		
	6	7.5	9
	-----tillers/ft ² -----		
10	20	16	13
12	24	19	16
14	28	22	19
16	32	26	21
18	36	29	24
20	40	32	27
22	44	35	29
24	48	38	32
26	52	42	35
28	56	45	37
30	60	48	40
32	64	51	43
34	68	54	45
36	72	58	48
38	76	61	51
40	80	64	53
42	84	67	56
44	88	70	59
46	92	74	61
48	96	77	64
50	100	80	67



Seeds per Head

Seeds per head is simply the product of the number of seeds per spikelet and the total number of spikelets per head. Oklahoma wheat averages between 2 grains and 2.5 grains per spikelet and from 8 spikelets to 12 spikelets per head for a total of 20 grains to 30 grains per head. This is the "typical" range and actual grains per head can deviate greatly from this range. Factors such as variety, emergence time, grazing, tiller population density, fertility, foliar disease, insect infestations, weed control and moisture availability will affect the final number of seeds per head.

Equation 1

$$\text{tillers per sq. ft.} = \frac{\text{tillers per ft. of row} \times 12}{\text{row spacing in inches}}$$

If a yield estimate is being calculated during or after grain fill, seeds per head can be estimated very precisely by counting seeds in random head samples from across the field. Estimates of seeds per head earlier in the year will be less reliable, as they rely upon several assumptions that may or may not prove to be correct as the crop year progresses. It is also important not to count seeds that are shriveled or light, as these will likely not make it to the combine tank and will not contribute much to yield if they do.

Seed Weight

The list of factors that can influence seeds per head also influence seed weight. Within a given environment, however, variety probably has the largest effect on seed size. Even though seed size influences final grain yield, it is important to note that small-seeded varieties frequently compensate

by producing more heads and more seeds per head than larger-seeded varieties. Given these caveats, most hard red winter wheat produced in Oklahoma will have between 13,000 and 16,000 seeds per pound. It is also important to note that seeds per pound will generally be larger in noncleaned wheat because small seeds are removed during the cleaning process.

Equation 2

$$\text{estimated grain yield (bu/A)} = \frac{\text{heads per sq. ft.} \times \text{grains per head} \times 726}{\text{seeds per pound}}$$

Calculating the estimate

Once estimates of the number of heads per square foot, seeds per head and seed weight have been established, they can be combined using Equation 2 to produce an estimated yield potential. The '726' value in Equation 2 is a constant that results from simplifying the equation. The results of Equation 2 using several possible combinations have been calculated in Table 2.

Table 2. Estimated yield potential for seeds per pound, seeds per head, and heads per square foot combinations likely to be found in Oklahoma wheat fields.

Heads per square foot	14,000 seeds per pound					15,000 seeds per pound					16,000 seeds per pound				
	Seeds per head					Seeds per head					Seeds per head				
	20	22	24	26	28	20	22	24	26	28	20	22	24	26	28
	grain yield estimate (bu/A)					grain yield estimate (bu/A)					grain yield estimate (bu/A)				
20	21	23	25	27	29	19	21	23	25	27	18	20	22	24	25
22	23	25	27	30	32	21	23	26	28	30	20	22	24	26	28
24	25	27	30	32	35	23	26	28	30	33	22	24	26	28	30
26	27	30	32	35	38	25	28	30	33	35	24	26	28	31	33
28	29	32	35	38	41	27	30	33	35	38	25	28	30	33	36
30	31	34	37	40	44	29	32	35	38	41	27	30	33	35	38
32	33	37	40	43	46	31	34	37	40	43	29	32	35	38	41
34	35	39	42	46	49	33	36	39	43	46	31	34	37	40	43
36	37	41	45	49	52	35	38	42	45	49	33	36	39	42	46
38	39	43	47	51	55	37	40	44	48	51	34	38	41	45	48
40	41	46	50	54	58	39	43	46	50	54	36	40	44	47	51
42	44	48	52	57	61	41	45	49	53	57	38	42	46	50	53
44	46	50	55	59	64	43	47	51	55	60	40	44	48	52	56
46	48	52	57	62	67	45	49	53	58	62	42	46	50	54	58
48	50	55	60	65	70	46	51	56	60	65	44	48	52	57	61
50	52	57	62	67	73	48	53	58	63	68	45	50	54	59	64
52	54	59	65	70	76	50	55	60	65	70	47	52	57	61	66
54	56	62	67	73	78	52	57	63	68	73	49	54	59	64	69
56	58	64	70	76	81	54	60	65	70	76	51	56	61	66	71
58	60	66	72	78	84	56	62	67	73	79	53	58	63	68	74
60	62	68	75	81	87	58	64	70	76	81	54	60	65	71	76
62	64	71	77	84	90	60	66	72	78	84	56	62	68	73	79
64	66	73	80	86	93	62	68	74	81	87	58	64	70	76	81
66	68	75	82	89	96	64	70	77	83	89	60	66	72	78	84
68	71	78	85	92	99	66	72	79	86	92	62	68	74	80	86
70	73	80	87	94	102	68	75	81	88	95	64	70	76	83	89

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Robert E. Whitson, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. 0309